










A polymorphism of *HMGA1* protects against proliferative diabetic retinopathy by impairing HMGA1-induced VEGFA expression

Eusebio Chiefari¹, Valeria Ventura¹, Carmelo Capula², Giorgio Randazzo³, Vincenzo Scorcia³, Monica Fedele⁴, Biagio Arcidiacono¹, Maria Teresa Nevolo¹, Francesco Luciano Bilotta¹, Michela Vitiello⁴, Camillo Palmieri³, Elio Gulletta¹, Alfredo Fusco^{4,5}, Daniela Foti¹, Raffaella Vero² & Antonio Brunetti^{1*}

Supplementary Information

Supplementary Figure



Supplementary Fig. S1: Full-length Western blots and gels. Western blots: Figure 1A, HMGA1, β-Tubulin; Figure 2A, HMGA1, Lamin A/C; Figure 2B (HepG2), HMGA1, Lamin A/C; Figure 2B (HUVEC), HMGA1, β-Tubulin; Figure 2C (left, ARPE-19), HMGA1, Lamin A/C; Figure 2C (right, ARPE-19), HMGA1, Lamin A/C; Figure 2D, VegfA, Hmga1. ChIP: Figure 3B, ChIP (HepG2) VEGFA; Figure 3B, ChIP (ARPE-19) VEGFA.