Nesting of colon and ovarian cancer cells in the endothelial niche is associated with alterations in glycan and lipid metabolism.

Anna Halama₁₊, Bella S Guerrouahen_{2,3,4+}, Jennifer Pasquier_{2,3}, Noothan J. Satheesh₁₊, Karsten Suhre _{1,5*} and Arash Rafii _{2,3,6*}.

†Contributed equally

- 1. Department of Physiology and Biophysics, Weill Cornell Medicine-Qatar, Qatar-Foundation, P.O. Box 24144, Doha, Qatar.
- 2. Stem Cell and Microenvironment Laboratory, Weill Cornell Medicine-Qatar, Education City, Qatar Foundation, Doha, Qatar.
- 3. Department of Genetic Medicine, Weill Cornell Medicine-Qatar, New York, NY 10065, USA.
- 4. Translational Medicine Division-Research Department, Sidra Medical and Research Center, PO Box 26999, Doha, Qatar.
- 5. Institute of Bioinformatics and Systems Biology, Helmholtz Zentrum München, German Research Center for Environmental Health, Neuherberg, Germany.
- 6. Department of Genetic Medicine and Obstetrics and Gynecology, Weill Cornell Medical College, Stem Cell and Microenvironment Laboratory, Weill Cornell Medical College in Qatar, Qatar-Foundation, P.O. Box 24144, Doha, Qatar.

Corresponding authors:

Dr. Karsten Suhre: Department of Physiology and Biophysics, Weill Cornell Medicine-Qatar, Doha, Qatar Weill Cornell Medical College in Qatar, Qatar-Foundation, P.O Box: 24144, Doha, Qatar. Phone: +974 4492 8482. Fax: +974 4492 8422. Email: kas2049@gatar-med.cornell.edu

Dr. Arash Rafii: Department of Genetic Medicine and Obstetrics and Gynecology, Weill Cornell Medicine-Qatar, Stem cell and microenvironment laboratory Weill Cornell Medical College in Qatar, Qatar-Foundation, P.O Box: 24144, Doha, Qatar. Phone: +974 3313 5828. Fax: +974 4492 8422. Email: jat2021@gatar-med.cornell.edu



