

Supplementary Information

Niclosamide reduces glucagon sensitivity via hepatic PKA inhibition in obese mice: Implications for glucose metabolism improvements in type 2 diabetes.

Md. Kamrul Hasan Chowdhury¹, Nigel Turner¹, Nicholas L Bentley¹, Abhirup Das¹, Lindsay E Wu¹, Dulama Richani², Sonia Bustamante³, Robert Gilchrist², Margaret J Morris¹, Peter R Shepherd⁴ and Greg C Smith^{1,*}

¹Department of Pharmacology, School of Medical Sciences, UNSW Australia, Sydney, Australia

²School of Women's & Children Health, UNSW Australia, Sydney, Australia

³Bioanalytical Mass Spectrometry Facility, UNSW Australia, Sydney, Australia

⁴Department of Molecular Medicine & Pathology, University of Auckland, Auckland, New Zealand

Correspondence


Greg C. Smith

Telephone: +61 (2) 93858075

Fax: +61 (2) 93850023

E-mail: g.smith@unsw.edu.au

Supplementary Figure S1



Supplementary Figure S1. NENS or Nic uncouples mitochondrial respiration. Oxygen consumption of isolated C57BL/6 hepatocytes using the Clark electrode with oligomycin and two forms of niclosamide (NENS or Nic; 500 nM and 5 μM). Each experiment is in duplicate and repeated three times independently. Statistical significance (P) was determined by multiple comparisons Two-way ANOVA test. The data is showed as the mean ± S.E.M. *p<0.01 or ***p<0.001.