

Supplementary Online Content

Gupta S, Halm EA, Rockey DC, et al. Comparative effectiveness of fecal immunochemical test outreach, colonoscopy outreach, and usual care for boosting colorectal cancer screening among the underserved: a randomized trial. *JAMA Intern Med.* doi:10.1001/jamainternmed.2013.9294.

eTable 1. Characteristics of Patients With CRC Detected

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable1. Characteristics of patients with CRC detected

GROUP	TNM STAGE	LOCATION	SCREEN DETECTED YES/NO*	DETECTED 2⁰ TO INTERVENTION YES/NO**
Colonoscopy Outreach	Stage I	Sigmoid Colon	Yes	Yes
	Stage I	Ascending Colon	No	No
FIT Outreach	Stage 0	Appendix	Yes	Yes
	Stage I	Cecum	Yes	Yes
	Stage IIIA	Splenic Flexure	Yes	Yes
	Stage 0	Ascending Colon	No	No
	Stage IV	Rectosigmoid Colon	No	No
	Unknown	Sigmoid Colon	No	No
Usual Care	Stage 0	Sigmoid Colon	Yes	No
	Stage 0	Sigmoid Colon	Yes	No
	Stage I	Ascending Colon	No	No
	Stage I	Rectosigmoid Colon	No	No
	Stage IIA	Rectum	No	No
	Stage IV	Splenic Flexure	No	No

*Refers to whether the colorectal cancer was diagnosed after presentation with signs/symptoms of cancer, or as a result of asymptomatic screening; **Refers to whether the cancer was detected as a result of asymptomatic screening that was initiated by our FIT or colonoscopy outreach interventions.