

RUNNING HEAD: SENSITIVITY TO COST OF LOW-NICOTINE CIGARETTES

**Supplementary Tables:**

Supplementary Table 1. Median demand parameters for Study Cigarettes.

		<b>Usual Brand</b>	<b>15.8</b>	<b>5.2</b>	<b>2.4</b>	<b>1.3</b>	<b>0.4</b>	<b>0.4 (HT)</b>
<b>Intensity</b>	Baseline	20	20	20	17	20	20	20
	Week 6	20	20	20	10	15	11.5	15
	Abstinence	20	20	17.5	10	10	10.5	10
<b>Omax</b>	Baseline	6.2	6.5	8	6	7.5	7.5	8
	Week 6	8	6	6	4	4	3	4
	Abstinence	6	7	5.55	2.45	3	2.7	2.85
<b>Pmax</b>	Baseline	0.5	0.5	0.55	0.5	0.5	0.5	0.7
	Week 6	0.4	0.4	0.4	0.3	0.3	0.3	0.3
	Abstinence	0.4	0.4	0.3	0.3	0.3	0.3	0.3
<b>Break point</b>	Baseline	0.9	1	1.5	0.9	2	0.5	2
	Week 6	0.7	0.8	0.6	0.45	0.55	1	0.5
	Abstinence	0.7	0.7	0.55	0.45	0.5	0.5	0.4
<b><math>\alpha</math></b>	Baseline	0.043	0.043	0.038	0.047	0.043	0.043	0.039
	Week 6	0.039	0.039	0.05	0.058	0.063	0.081	0.062
	Abstinence	0.036	0.031	0.046	0.054	0.065	0.085	0.066

RUNNING HEAD: SENSITIVITY TO COST OF LOW-NICOTINE CIGARETTES

Supplementary Table 2. Median demand parameters for Usual Brand Cigarettes.

		<b>Usual Brand</b>	<b>15.8</b>	<b>5.2</b>	<b>2.4</b>	<b>1.3</b>	<b>0.4</b>	<b>0.4 (HT)</b>
<b>Inten sity</b>	Baseline	20	20	20	17	20	20	20
	Week 6	20	20	20	18	20	15.5	20
	Abstinence	20	20	20	15	15	15	16.5
<b>Omax</b>	Baseline	6.2	6.5	8	6	7.5	7.5	8
	Week 6	8	6.8	7.35	7	6	6	6
	Abstinence	6	7.5	6	6	4.65	6	4.75
<b>Pmax</b>	Baseline	0.5	0.5	0.55	0.5	0.5	0.5	0.7
	Week 6	0.4	0.4	0.5	0.5	0.4	0.45	0.4
	Abstinence	0.4	0.4	0.4	0.4	0.4	0.4	0.4
<b>Break point</b>	Baseline	0.9	1	1.5	0.9	2	1	2
	Week 6	0.7	0.8	0.8	0.8	0.8	0.9	0.9
	Abstinence	0.7	0.8	0.7	0.8	0.7	0.8	0.7
<b><math>\alpha</math></b>	Baseline	0.043	0.043	0.038	0.047	0.043	0.043	0.039
	Week 6	0.039	0.041	0.036	0.035	0.045	0.042	0.042
	Abstinence	0.036	0.032	0.041	0.04	0.048	0.048	0.047