

Neutrophil gelatinase-associated lipocalin (NGAL) fails as an early predictor of contrast induced nephropathy in chronic kidney disease (ANTI-CI-AKI study)

Werner Ribitsch¹, Gernot Schilcher^{1,2*}, Franz Quehenberger³, Stefan Pilz⁴, Rupert H Portugaller⁵, Martini
Truschnig-Wilders⁶, Robert Zweiker⁷, Marianne Brodmann⁸, Philipp Stiegler⁹, Alexander R Rosenkranz¹, John
W Pickering¹⁰ and Joerg H Horina¹

¹Clinical Division of Nephrology, Department of Internal Medicine, Medical University of Graz (MUG), Austria

²Intensive Care Unit, Department of Internal Medicine, MUG, Austria

³Institute for Medical Informatics, Statistics and Documentation, Medical University of Graz, Austria

⁴Division of Endocrinology and Metabolism, Department of Internal Medicine,
Medical University of Graz, Austria

⁵Department of Vascular and Interventional Radiology, University Clinic of Radiology,
Medical University of Graz, Austria

⁶Clinical Institute for Medical and Chemical Laboratory Diagnostics, Medical University of Graz, Austria

⁷Division of Cardiology, Department of Internal Medicine, Medical University of Graz, Austria

⁸Division of Angiology, Department of Internal Medicine, Medical University of Graz, Austria

⁹Division of Transplantation Surgery, Medical University of Graz, Austria

¹⁰Department of Medicine, University of Otago Christchurch and Emergency Medicine Department,
Christchurch Hospital, Christchurch, New Zealand

Supplementary Appendix

Supplemental Table 1. Adverse events (AE) during the ANTI-CIN -study

Adverse event (n)	AE Day -1	SAE Day-1	AE Day 0	SAE Day 0	AE Day 1	SAE Day 1	AE Day 2	SAE Day 2
Pain (<i>localization</i>)								
Headache	7		8		6		2	
Musculoskeletal system	20		69		25		20	1
Thorax	1		4	3	1			
Abdomen	1		4		1		1	
Cardiac (Arrhythmias, NSTEMI, pulmonary edema, hypotension)		4	27	6	7	4	1	
Hypertensive crisis	8		45		9			
Neurological (Dizziness, mental confusion,	6	1	5	4	11	1	2	2
Nausea, vomiting	2	3	11		6		3	
Dyspnoea			7		6		2	
Epistaxis	1	1						
Infection	1	4	2	1	4		2	2
Dermatological (rash, exanthema, pruritus)	1		16		10			1
Peri-interventional complications								
Dissection								
Bradycardia			5	30				
Tachycardia			3					
Hypotension			5					
Hypertension			19					
Hematoma			10					
Embolism			2					
Death				1				
Other			13					
Total	48	13	255	45	86	5	33	6

Urinary NGAL and contrast induced acute kidney injury

Supplemental Table 2. Adverse reactions (AR) during the ANTI-CIN -study

Adverse reaction (n)	AR Day -1	SAR Day-1	AR Day 0
Headache	1		
Cardial (Arrhythmias, NSTEMI, pulmonary edema, hypotension)	1		
Hypertensive crisis		2	1
Dyspnoea	6		
Neurological	1		
<i>Total</i>	9	2	1