

**Supplemental Table 4. Cortisol Levels for Individuals During Adulthood**

<b>Subjects</b>	<b>Baseline / Non-stress</b>		<b>Human Intruder Paradigm</b>		<b>Diurnal Rhythm</b>		
	<b>Lights-On</b>	<b>+30min</b>	<b>Pre-stress</b>	<b>Post-stress</b>	<b>Lights-On</b>	<b>Mid-day</b>	<b>Lights-Off</b>
Neo-C-1	25.74	22.72	20.52	24.55	19.32	7.79	2.50
Neo-C-2	29.59	29.48	27.93	32.58	21.25	11.55	4.88
Neo-C-3	30.20	23.30	23.80	32.54	23.09	7.92	2.50
Neo-C-4	21.31	16.18	17.53	28.61	22.59	7.35	2.50
Neo-C-5	28.34	22.37	26.87	35.84	23.08	10.57	2.50
Neo-C-7	31.56	27.58	24.48	41.08	21.54	10.18	3.52
Neo-Hibo-1	21.25	18.83	25.22	27.26	19.13	9.72	9.26
Neo-Hibo-2	14.30	13.42	18.22	14.15	15.76	9.80	2.50
Neo-Hibo-3	15.14	14.85	21.15	14.84	18.71	16.27	3.09
Neo-Hibo-4	25.26	27.06	24.51	30.78	15.30	9.69	2.50
Neo-Hibo-5	27.52	22.30	28.91	33.58	22.47	15.74	4.95
Neo-Hibo-6	18.13	9.63	17.69	21.73	15.70	8.60	3.98

Cortisol levels ( $\mu\text{g}/\text{dl}$ ) for all subjects during adulthood (6-8 years). Cortisol was measured by Diagnostic Systems Laboratories (DSL) radioimmunoassay for Baseline/Non-stress and Human Intruder Paradigm days, whereas Diurnal Rhythm of cortisol was measured using liquid chromatography - mass spectrometry (LC-MS). Neo-C-7 has also been reported as Neo-C-11 in other publications (Weiss & Bachevalier, 2016; Weiss et al, 2016).