

SUPPLEMENTARY FILE

Hepatic cyst infection during use of the somatostatin analogue lanreotide in autosomal dominant polycystic kidney disease: an interim analysis of the multicenter, open label, randomized DIPAK-1 study

Drug Safety

Marten A Lantinga^{1*}, MD; Hedwig M.A. D'Agnolo^{1*}, MD; Niek F. Castelein², MD; Johan W. de Fijter³, MD, PhD; Esther Meijer², MD, PhD; Annemarie L. Messchendorp², MD; Dorien J.M. Peters⁴, PhD; Mahdi Salih⁵, MD; Edwin M. Spithoven², MD PhD; Darius Soonawala³, MD; Folkert W. Visser², MD PhD; Jack F. Wetzels⁶, MD PhD; Bob Zietse⁵, MD PhD; Joost P.H. Drenth¹, MD, PhD, and Ron T. Gansevoort², MD PhD; on behalf of the DIPAK Consortium

*both authors contributed equally.

¹ Dept. of Gastroenterology and Hepatology, Radboud University Medical Center Nijmegen, the Netherlands

² Dept. of Nephrology, University Medical Center Groningen, the Netherlands

³ Dept. of Nephrology, Leiden University Medical Center, the Netherlands

⁴ Dept. of Human Genetics, Leiden University Medical Center, Leiden, the Netherlands

⁵ Dept. of Nephrology, Erasmus Medical Center Rotterdam, the Netherlands

⁶ Dept. of Nephrology, Radboud University Medical Center Nijmegen, the Netherlands

Corresponding author: Ron T. Gansevoort, MD PhD: r.t.gansevoort@umcg.nl

Content: Supplement Figure 1 and Legend Supplement Figure 1

Figure 1. ^{18}F -FDG PET/CT results showing signs of hepatic cyst infection in case 1(A) and case 5 (B)

