

**Supplemental Table 1. Primers sequences used for this study**

<b>Gene</b>	<b>Forward Sequence (5'-3')</b>	<b>Reverse Sequence (5'-3')</b>
<i>BRD2</i>	CTACGTAAGAAACCCCGGAAG	GCTTTTTCTCCAAAGCCAGTT
<i>BRD3</i>	CCTCAGGGAGATGCTATCCA	ATGTCGTGGTAGTCGTGCAG
<i>BRD4</i>	TTTGAGACCCTGAAGCCGTC	TTAGGCAGGACCTGTTTCGG
<i>JAG1</i>	CTATGATGAGGGGGATGCT	CGTCCATT CAGGCACTGG
<i>ACTB</i>	ATTGGCAATGAGCGGTTCC	GGTAGTTTCGTGGATGCCACA
<i>POLR2A</i>	ACTTTTGGTACATTGTGGCTTCAA	CCGCCAGGACAAACCAGTAT

The Supplement Table 1 lists the qPCR primers used in this study.

**Supplemental Table 2. ChIP primers sequences used for this study**

<b>Region</b>	<b>Forward Sequence (5'-3')</b>	<b>Reverse Sequence (5'-3')</b>
Jagged1 proximal promoter	AGCAGCGTCTGCCAGGT	TGACTGAGCGGTCCGAGC
Jagged1 distal promoter	GGAGACTGGTGTGATGGA	GAGCAAGTTTACAGCTGGT

The Supplement Table 2 lists the ChIP primers used in this study.