Supplemental Table 2: Bivariate comparisons of individuals who did or did not experience readmission for fall-related injury.

## 90 Day Fall

	Test Statistic	p value
Private Payer	<i>X</i> <sup>2</sup> = 1.189e-56	p < .001
Male	$X^2 = 0.2911$	0.2911
White	<i>X</i> <sup>2</sup> = 7.61e-16	p < .001
Admission Psychiatric Dx	$X^2 = 0.0001597$	p < .001
Admission via ED	<i>X</i> <sup>2</sup> = 1.537e-18	p < .001
Total medications	<i>t</i> = 4.072e-10	p < .001
Age	<i>t</i> = 0.0002198	p < .001
Charlson	<i>t</i> = 4.009e-38	p < .001
Burden Score	<i>t</i> = 3.232e-09	p < .001

## 180 Day Fall

	Test Statistic	p value
Private Payer	<i>X</i> <sup>2</sup> = 6.147e-108	p < .001
Male	$X^2 = 0.1702$	0.1702
White	$X^2 = 6.574e-33$	p < .001
Admission Psychiatric Dx	$X^2 = 2.98e-07$	p < .001
Admission via ED	<i>X</i> <sup>2</sup> = 2.952e-31	p < .001
Total medications	<i>t</i> = 4.489e-18	p < .001

	Age	<i>t</i> = 3.102e-07	p < .001
	Charlson	<i>t</i> = 5.797e-78	p < .001
	Burden Score	<i>t</i> = 7.067e-18	p < .001