

Table S1. Summary of injection sessions

Cpf1 (ng/μl)	Target A (ng/μl)	Target B (ng/μl)	Egg strain (♀ x ♂)	# eggs implanted	# pups or embryos recovered	# founders targeted	% founders targeted
10	2.5	2.5	B6xFVB	128	24	5/24	20.8%
10	2.5	2.5	B6xFVB	101	30	1/30	3.3%
10	2.5	2.5	FVBxB6	180	54	1/54	1.9%
10	2.5	2.5	FVBxB6	110	17	0/17	0.0%
10	2.5	2.5	FVBxB6	168	36	0/36	0.0%
10	2.5	-	FVBxB6	40	1	0/1	0.0%
10	2.5	-	FVBxB6	144	45	0/45	0.0%
50	100	100	B6xFVB	50	21	18/21	85.7%
50	100	100	B6xFVB	48	23	21/23	91.3%
50	100	100	B6 x B6	62	16	13/16	81.2%
50	100	100	FVBxB6	80	2	2/2	100%
50	100	-	FVBxB6	72	11	5/11	45.5%
50	100	-	FVBxB6	100	17	7/17	41.2%
Average across all 12 injection sessions:						73/297	24.6%
Average for 7 low concentration injection sessions:						7/207	3.4%
Average for 6 high concentration injection sessions:						66/90	73.3%