

Supplementary Information

High expression of AKR1B10 predicts low risk of early tumor recurrence in patients with hepatitis B virus-related hepatocellular carcinoma

Yan-Yan Wang^{1,3,*}, Lu-Nan Qi^{1,2,3,*}, Jian-Hong Zhong^{1,2,3}, Hong-Gui Qin^{1,3}, Jia-Zhou Ye^{1,2,3}, Shi-Dong Lu^{1,3}, Liang Ma^{1,2,3}, Xue-Mei You^{1,2,3}, Bang-De Xiang^{1,2,3}, and Le-Qun Li^{1,2,3}

1 Department of Hepatobiliary Surgery, Affiliated Tumor Hospital of Guangxi Medical University, Nanning 530021, PR China

2 Guangxi Cancer Institute, Nanning 530021, PR China

3 Key Laboratory of Early Prevention and Treatment of Regional High-Incidence-Tumors, Ministry of Education, Nanning 530021, PR China

*These authors contributed equally to this work.

Correspondence to: Dr Le-Qun Li (email: wyy857355056@163.com) or Xue-Mei You (email: 397845319@qq.com), Department of Hepatobiliary Surgery, Affiliated Tumor Hospital of Guangxi Medical University, He Di Rd. #71, Nanning 530021, China. Phone: +86-771-5330855. Fax: +86-771-5312000.

Manuscript type: original article.

SUPPLEMENTARY FIGURES

Supplementary Figure 1. Full-length blots of AKR1B10 (A) and GAPDH (B)

detected by Western blotting assay.

