

Supplementary Table 1. Primer sequences for qRT-PCR

Gene Symbol	Forward Primer	Reverse Primer
B2M [1]	TTTCATCCATCCGACATTG	CGGCAGGCATACTCATCTTT
ARRDC3	GCGAAAGTACGCTGGACTGA	TGAGGTAGCGAGTGGTGTCT
CA12	ACTTATTTTGGTCCTGATGGGG	TGGCCATTGTTGGTCAGGAG
CDKN3	CTTGTCTTGTAGCTGCTTGTCTC	TGCAGCTAATTTGTCCCGAAAC
FAM189A2 [2]	CTTGAGGACCAGGTCGAAGAGT	CTCCTTTCAGTCTGTGTTGCAGC
MME [2]	CTTTAGTGCCAGCAGTCCAAC	CACCAGTCAACGAGGTCTCCAT
PLXNA4 [2]	TCGTGCGGATTGAGCCAGAATG	TGATGTGCTCCTTCCCTCCATG
PPARG	CGACCAGCTGAATCCAGAGT	GATGCGGATGGCCACCTCTT
PTP4A3	ACAAGCACAGGGATCTCGTT	CTTCAGGTCCTCAATGAAGGTG
RALGPS2	GCAGTCAGTCCTCTGTTGCT	TCTGACCCGCATATTCTTCTGG
SNX33	GCTTTGTGCGTTCTGGAGTG	TGGTCTGTTTTGTGGGGTCC
TM4SF18	ATGTTGCCAGAGTGAAAAGTGC	CTGTAAGGAAACGTCCAGCA
TMEM121 [2]	AGCACATAGCGCCGAGAAGAT	CGTTTTTGCCGACGAAGATGGC
ZNF292	GTGTGAAGATGGCGGACGAA	GCATATTCTAGGAGTGTCTGGCA

References

1. Weijers EM, van Wijhe MH, Joosten L, et al. (2010) Molecular weight fibrinogen variants alter gene expression and functional characteristics of human endothelial cells. *J Thromb Haemost* 8:2800–9. doi: 10.1111/j.1538-7836.2010.04096.x
2. OriGene. <http://www.origene.com/>. Accessed 29 Jan 2016