

Supplementary Table 1. Univariate and multivariate Cox proportional hazard models for overall survival

	Overall survival		1-year survival	
	Univariate HR (95% CI)	Multivariate HR* (95% CI)	Univariate HR (95% CI)	Multivariate HR* (95% CI)
D-dimer quartiles (range, µg/mL)				
Q1 (< 2.08)	(Ref)	(Ref)		
Q2 (2.08–9.06)	2.01 (1.35–2.99) [‡]	1.44 (0.92–2.26)	2.30 (1.45–3.67) [‡]	1.62 (0.95–2.76)
Q3–Q4 (> 9.06)	3.08 (2.15–4.40) [‡]	2.19 (1.46–3.31) [‡]	3.78 (2.49–5.75) [‡]	2.70 (1.68–4.35) [‡]
Systemic metastasis	3.51 (2.51–4.91) [‡]	2.91 (1.95–4.34) [‡]	3.48 (2.39–5.07) [‡]	2.96 (1.86–4.70) [‡]
Stroke mechanism [†]	1.75 (1.30–2.37) [‡]		1.99 (1.41–2.81) [‡]	
Age	1.00 (0.99–1.01)		1.00 (0.98–1.01)	
NIHSS	1.03 (1.01–1.06) [‡]	1.03 (1.01–1.06) [‡]	1.04 (1.01–1.06) [‡]	1.03 (1.00–1.06) [‡]
Hepatobiliary/pancreatic cancer	2.01 (1.45–2.77) [‡]	2.25 (1.54–3.29) [‡]	2.08 (1.49–2.90) [‡]	2.23 (1.51–3.28) [‡]
Adenocarcinoma	1.13 (0.84–1.52)		0.87 (0.63–1.20)	0.68 (0.47–1.00) [‡]
Atrial fibrillation	0.91 (0.57–1.45)		0.83 (0.49–1.41)	

*Multivariate models adjusted for variables used in univariate analysis with a forward stepwise method; [†]Cryptogenic vs conventional mechanisms (reference); [‡]*P* < 0.05.

Q1, Q2, Q3, and Q4 denote quartiles.