

Supplementary Table 1. Clinical characteristics and outcome of patients with and without recanalisation therapy (only patients with available follow-up imaging)

	Patients without recanalization therapy (N=129)	Patients with recanalization therapy (N=32)	P value
Age (year)	77.0 (\pm 7.9)	75.7 (\pm 11.7)	0.54
Women	64 (49.6)	17 (53.1)	0.84
NOAC			0.78
Apixaban	20 (15.5)	5 (15.6)	
Dabigatran	36 (27.9)	7 (21.9)	
Rivaroxaban	73 (56.6)	20 (62.5)	
Concomitant platelet inhibition	13 (10.1)	3 (9.4)	> 0.99
CHA ₂ DS ₂ -VASc score	5 (3–6)	4 (3–5)	0.02
HAS-BLED score	4 (3–4)	3 (3–3)	< 0.001
Medical history			
Ischemic stroke/TIA	60 (46.5)	6 (18.8)	0.005
Intracranial haemorrhage	3 (2.3)	0 (0)	> 0.99
Atrial fibrillation	119 (92.2)	26 (81.3)	0.09
Hypertension	106 (82.2)	26 (81.3)	> 0.99
Hyperlipidemia	47 (36.4)	7 (21.9)	0.15
Diabetes mellitus	44 (34.1)	4 (12.5)	0.02
Heart failure	34 (26.4)	10 (31.3)	0.66
Peripheral vascular disease	9 (7.0)	1 (3.1)	0.69
Renal function			
GFR < 60 mL/min	53 (41.1)	10 (31.3)	0.30
Creatinin level (mg/dl)	1.01 (0.87–1.30)	0.9 (0.79–1.08)	0.02
NIHSS at admission	4 (2–7)	19 (14–22)	< 0.001
Modified Rankin scale score			
Pre stroke	1 (0–2)	1 (0–2)	0.24
At admission	3 (1–4)	5 (5–5)	< 0.001
At discharge	2 (1–4)	4 (3–5)	< 0.001
At 3-months*	2 (1–4)	4 (3–6)	< 0.001
mRS 0–2 at 3 months	66 (51.2)	5 (15.6)	< 0.001
In-hospital mortality	4 (3.1)	3 (9.4)	0.14
Time since last intake NOAC (hour)	9 (4–18)	10 (5–24)	0.45
Admission imaging			
ASPECTS	10 (9–10)	8 (7–10)	< 0.001
Haemorrhagic transformation			0.54
HI1	2	0	
HI2	3	0	
PH1	0	0	
PH2	0	0	
PHr1/PHr2	0/0	0/0	
Other haemorrhage			–
SAH	0	0	
IVH	0	0	
sICH	0	0	–
Follow-up imaging until day 7			
ASPECTS	9 (8–10)	6 (3–8)	< 0.001
Infarct size			< 0.001
Small/medium	99 (76.7)	10 (31.3)	
Large anterior/posterior	30 (23.3)	22 (68.8)	
Haemorrhagic transformation	18 (14.0)	13 (40.6)	0.002
Haemorrhagic transformation			0.03
HI1	10	5	
HI 2	2	3	
PH1	0	5	
PH2	1	1	
PHr1 / PHr2	0 / 0	0 / 0	

(Continued to the next page)

Supplementary Table 1. Continued from the previous page

	Patients without recanalization therapy (N = 129)	Patients with recanalization therapy (N = 32)	P value
Other haemorrhage			
SAH	0	2	0.04
IVH	0	1	-
sICH	1	1	-

Data are mean (\pm SD), median (IQR) or n (%).

*3-months modified Rankin data available for n = 117/129 (90.7%, patients without recanalization therapy), and n = 31/32 patients (96.9%, patients with re-canälization therapy).