

S1 Table. PCR primers used to amplify gene-specific cDNAs for the *in vitro* transcription of digoxigenin-labeled RNA probes.

Gene Symbol	Primer Sequences (5' to 3')
<i>aldh1a2</i>	F: ACTGCCAGGAGAGGTGAAGA R: CG TAATACGACTCACTATAGGG GCCTGTTCTAATGCCAGCTC
<i>bcl2l10</i>	F: GCACAGAACCAAATTCCGAT R: CG TAATACGACTCACTATAGGG GAAATCGCTTGGAACCAAAA
<i>cd44</i>	F: GTCTGCATTGCTGTCCCTCT R: CG TAATACGACTCACTATAGGG TGCTTCTGTAGGGCTGTGTG
<i>dlx5a</i>	F: AGAGACGCTTGACGCTTGAC R: CG TAATACGACTCACTATAGGG AACGTGCCGGGTGCCTGAAG
<i>fgf20a</i>	F: ATGGGTGCAGTCGGCGAGC R: CG TAATACGACTCACTATAGGG AAGCTCAGGAACTCGCTCTGGA
<i>fn1b</i>	F: TCGTGGACTGGAGCCAGGCAT R: CG TAATACGACTCACTATAGGG ACGCAATGCTTCCACCCGCA
<i>has1</i>	F: TGTGTCTCTGGTCCGAGCCTT R: CG TAATACGACTCACTATAGGG TCCCTCTTTCCTCCCCATTTCTGC
<i>has2</i>	F: ACTTCAACCTGTGCGCTTGGGG R: CG TAATACGACTCACTATAGGG GCCGGGTCCAGCATGGTGTC
<i>has3</i>	F: CACACAGAGCAGCACCATCT R: CG TAATACGACTCACTATAGGG GATTGAGCCAACGCAGGTAT
<i>hmmr</i>	F: CATGGAGCGCCGGGTGTCTG R: CG TAATACGACTCACTATAGGG GCTTCTGCTGGCCCTGCTGT
<i>hyal2</i>	F: AGAGTGGCGGCCCTTTGGA R: CG TAATACGACTCACTATAGGG CGCCGCTCGATTCCCGTTGT
<i>hyal3</i>	F: CCGCCTGGCCCTCTACCCTT R: CG TAATACGACTCACTATAGGG GCAGTTCGTCCTCCGGCCC
<i>hyal4</i>	F: GCACAACTGTTCCAAAGTGAAGGCA R: CG TAATACGACTCACTATAGGG TGATGTGAGTCTGGGGTGGGGT
<i>hyal6</i>	F: AAAGTGGCGACCGCAGTGGG R: CG TAATACGACTCACTATAGGG AGGCTCTGCCATCTGGGGCA
<i>junba</i>	F: ACGGAGAGAACTTGCGGACT R: CG TAATACGACTCACTATAGGG CTGCTCTTCCAGTGTTTCCC
<i>junbb</i>	F: ACGTGACCGAGCCGCCCTAT R: CG TAATACGACTCACTATAGGG GGGCCGCAGTGCCGTTCTC
<i>mmp9</i>	F: CCCTTGGCTCTGGACCAGCCAT R: CG TAATACGACTCACTATAGGG AGTCGCTTGGTAGGCCGAGCT
<i>snx18a</i>	F: CCCTCTGGCTACAGCATCTC R: CG TAATACGACTCACTATAGGG AGTTGGTGAGGTGTCCTTGG
<i>socs3b</i>	F: GCTCCCATCTAGCCCTCTCT R: CG TAATACGACTCACTATAGGG CGGTCATCAGACCAGACCTT
<i>vcana</i>	F: TGAAAAGTGGAGACCGAACC R: CG TAATACGACTCACTATAGGG AGGAAATCCTGGGAGAGTCC

The T7 promoter sequence is indicated in blue.