

Supplementary Table S4. Elimination processes.

Renal clearance processes considered in the developed PBPK models of APAP and CAF. Kidney plasma clearances were parameterized in order to match experimental observations.

Drug / Metabolite	Plasma clearance [l/min/kg]	Ref.
APAP	1.27E-04	(1,2)
APAPC	4.38E-04	(1)
APAPG	1.19E-03	(1)
APAPS	4.56E-03	(1)
NAPQI	4.62E-03	adjusted from (3)
CAF	3.00E-03	(4)

REFERENCES

1. Critchley J a, Nimmo GR, Gregson C a, Woolhouse NM, Prescott LF. Inter-subject and ethnic differences in paracetamol metabolism. *Br J Clin Pharmacol.* 1986;22(6):649–57.
2. Wishart DS, Knox C, Guo AC, Shrivastava S, Hassanali M, Stothard P, et al. DrugBank: a comprehensive resource for in silico drug discovery and exploration. *Nucleic Acids Res.* 2006 Jan 1;34(Database issue):D668–72.
3. Krauss M, Schaller S, Borchers S, Findeisen R, Lippert J, Kuepfer L. Integrating Cellular Metabolism into a Multiscale Whole-Body Model. Papin JA, editor. *PLoS Comput Biol.* 2012 Oct 25;8(10):e1002750.
4. Tang-Liu DD, Williams RL, Riegelman S. Disposition of caffeine and its metabolites in man. *J Pharmacol Exp Ther.* 1983 Jan;224(1):180–5.