

## Supplementary Materials: Metabolomic Biomarkers in Urine of Cushing's Syndrome Patients

Alicja Kotłowska, Tomasz Puzyn, Krzysztof Sworczak, Piotr Stepnowski and Piotr Szefer

**Table S1.** Metadata obtained for patients. All the results apart from gender are expressed as medians (5th, 95th percentile).

<b>Metadata</b>	<b>Adrenal Incidentaloma (n = 25)</b>	<b>Cushing's Syndrome (n = 16)</b>
Age (years)	49 (43, 57)	51 (44, 58)
Gender (%)	72% females 28% males	68.75% females 31.25% males
Height (cm)	163 (157, 176)	167.5 (158, 178)
Weight (kg)	77 (71, 89)	86 (74.5, 101)
BMI (kg/m <sup>2</sup> )	28.9 (27.9, 30.3)	30.8 (29.9, 33)
UFC (nmol/24 h) <sup>1</sup>	112 (86, 144)	905 (690, 1409)
Serum cortisol after DXM suppression test (nmol/L) <sup>2</sup>	66 (36, 85)	279 (170, 367)

<sup>1</sup> UFC normal range: 12–330 nmol/24 h, values > 330 nmol/24 h indicated hypercortisolism; <sup>2</sup> Serum cortisol after dexamethasone (DXM) suppression test: concentrations < 50 nmol/L excluded hypercortisolism, intermediate cut-off point of 94 nmol/L indicated subclinical hypercortisolism, >140 nmol/L indicated hypercortisolism.