

Ibuprofen results in alterations of human fetal testis development

Millissia Ben Maamar¹, Laurianne Lesné¹, Kristin Hennig², Christèle Desdoits-Lethimonier¹, Karen R. Kilcoyne³, Isabelle Coiffec¹, Antoine D. Rolland¹, Cécile Chevrier¹, David M. Kristensen⁴, Vincent Lavoué⁵, Jean-Philippe Antignac², Bruno Le Bizec², Nathalie Dejucq-Rainsford¹, Rod T. Mitchell³, Séverine Mazaud-Guittot¹, Bernard Jégou^{1,6*}

¹Institut national de la santé et de la recherche médicale (Inserm), Institut de recherche en santé, environnement et travail (Irset – Inserm UMR 1085), 9 Avenue Léon Bernard, F-35000 RENNES, France. Université de Rennes 1, F-35043 RENNES, France.

²LUNAM Université, Oniris, USC INRA 1329, Laboratoire d'Etude des Résidus et Contaminants dans les Aliments (LABERCA), Nantes, F-44307, France.

³MRC Centre for Reproductive Health, University of Edinburgh, Queens Medical Research Institute, 47 Little France Crescent, Edinburgh, EH16 4TJ.

⁴Laboratorium of Genomic and Molecular Biomedicine, Department of Biology, University of Copenhagen, Ole Maaløes Vej 5, DK-2200 Copenhagen N, Denmark.

⁵CHU de Rennes, Service de Gynécologie, Hôpital Sud, 16, boulevard de Bulgarie, F-35700 Rennes, France; Université de Rennes 1, Faculté de Médecine, F-35043 RENNES, France.

⁶Ecole des hautes études en santé publique (EHESP), Avenue Léon Bernard, F-35043 RENNES, France.

*To whom correspondence should be addressed: bernard.jegou@inserm.fr

Figure S1.




Figure S1. Body weight (A), Grafts retrieved (B) and Total graft weight (C) for individual host mice carrying human fetal testis xenografts (14-17 GW; n=4) exposed for 7d to vehicle (corn oil; open circles) or ibuprofen (10 mg/kg 3 times daily; closed circles); with overall mean \pm SEM for vehicle (white bars) and ibuprofen (black bars). (D) Plasma ibuprofen concentration in vehicle (open circles) and ibuprofen (closed circles) exposed host mice. Data analyzed by two-way ANOVA.