

Supplementary Material

Combination of Endothelial-Monocyte-Activating Polypeptide-II with Temozolomide Suppress Malignant Biological Behaviors of Human Glioblastoma Stem Cells via miR-590-3p/MACC1 Inhibiting PI3K/AKT/mTOR Signal Pathway

Wei Zhou^{1,2}, Libo Liu^{3,4}, Yixue Xue^{3,4}, Jian Zheng^{1,2}, Xiaobai Liu^{1,2}, Jun Ma^{3,4}, Zhen Li^{1,2}, Yunhui Liu^{1,2*}

¹Department of Neurosurgery, Shengjing Hospital of China Medical University, Shenyang China

²Liaoning Research Center for Translational Medicine in Nervous System Disease, Shenyang China

³Department of Neurobiology, College of Basic Medicine, China Medical University, Shenyang China

⁴ Key Laboratory of Cell Biology, Ministry of Public Health of China, and Key Laboratory of Medical Cell Biology, Ministry of Education of China, China Medical University, Shenyang China

*Correspondence: Yunhui Liu: liuyh_cmuns@163.com

1. Supplementary data

Supplementary Table S1. The clinically-relevant details of patient tissues.

The study was approved by the Ethics Committee of Shengjing Hospital of China Medical University. The grade was identified according to the WHO classification by experienced clinical pathologists.

Patient number	Age	Sex	WHO Grade	Histology type	Lobe
1	32	M		Craniocerebral trauma	Frontal lobe
2	22	M		Craniocerebral trauma	Frontal lobe
3	38	F		Craniocerebral trauma	Frontal lobe
4	47	F	I	Pilocytic astrocytoma	Temporal lobe
5	59	F	I	Pilocytic astrocytoma	Temporal lobe
6	45	M	II	Diffuse astrocytoma	Temporal lobe
7	61	M	IV	Glioblastoma	Frontal lobe
8	69	F	IV	Glioblastoma	Frontal lobe
9	59	M	IV	Glioblastoma	Frontal lobe

