

Cancer-associated fibroblasts promote an immunosuppressive microenvironment through the induction and accumulation of protumoral macrophages

SUPPLEMENTARY TABLE

Supplementary Table S1: List of primers used for quantitative RT-PCR in this study

Gene	Forward primer	Reverse primer
<i>ARG1</i>	5'-AAAGGCTGGTCTGCTTGAGAA-3'	5'-GTCATTAGGGATGTCAGCAAAGG-3'
<i>IL10</i>	5'-GAGATGCCTTCAGCAGAGTGAAGA-3'	5'-AGGCTTGGCAACCCAGGTAAC-3'
<i>TGFBI</i>	5'-AGCGACTCGCCAGAGTGGTTA-3'	5'-GCAGTGTGTTATCCCTGCTGTCA-3'
<i>VEGFA</i>	5'-ACTTCCCAAATCACTGTGG-3'	5'-GTCACTCACTTTGCCCTGT-3'
<i>TNF</i>	5'-TGCTTGTTCCCTCAGCCTCTT-3'	5'-CAGAGGGCTGATTAGAGAGAGGT-3'
<i>IL1B</i>	5'-TCGCCAGTCAAATGATGGCTTA-3'	5'-GTCCATGGCCACAACAACACTGA-3'
<i>GAPDH</i>	5'-GCACCGTCAAGGCTGAGAAC-3'	5'-ATGGTGGTGAAGACGCCAGT-3'