

Table 1 supplementary.

Multivariate logistic analysis with HCV RNA as a continuous variable.

	<i>Rapid viral response</i>			<i>Undetectable or VL\leq15 UI</i>		
	<i>OR</i>	<i>CI 95%</i>	<i>P</i>	<i>OR</i>	<i>CI 95%</i>	<i>P</i>
Treatment initiation (Season A vs. Season B)	1.93	1.22-3.07	0.005	1.81	1.08-3.04	0.025
Type of therapy (lead-in no vs. yes)	23.41	10.57-51.86	<0.001	67.94	37.76-122.25	<0.001
HCV RNA (IU ml ⁻¹)	1.00	1.00-1.00	0.712	1.00	1.00-1.00	0.181
IL-28 polymorphism (CC vs. no CC)	1.85	1.09-3.12	0.021	2.09	1.11-3.93	0.021
Naïve vs. experienced	1.49	0.89-2.52	0.128	0.70	0.39-1.23	0.215
Genotype (1b vs. 1a)	1.38	0.76-2.48	0.284	0.79	0.42-1.48	0.471
Diabetes Mellitus (no vs. yes)	1.30	0.64-2.64	0.461	1.52	0.72-3.18	0.272
Age (\geq 50 vs. < 50 years)	0.94	0.57-1.54	0.806	1.06	0.61-1.84	0.837
Fibrosis (\geq F3 vs. <F3)	0.91	0.51-1.64	0.762	1.62	0.80-3.26	0.004

Table 2 supplementary. Multivariate logistic analysis including relapsers in the naive group.

	<i>Rapid viral response</i>			<i>Undetectable or VL≤15 UI</i>		
	<i>OR</i>	<i>CI 95%</i>	<i>P</i>	<i>OR</i>	<i>CI 95%</i>	<i>P</i>
Treatment initiation (Season A vs. Season B)	1.96	1.23-3.12	<0.001	1.74	1.03-2.94	0.038
Type of therapy (lead-in no vs. yes)	24.16	10.85-53.75	0.004	64.76	35.99-116.56	<0.001
HCV RNA (IU ml ⁻¹)	1.00	1.00-1.00	0.713	1.00	1.00-1.00	0.263
IL-28 polymorphism (CC vs. no CC)	1.81	1.07-3.04	0.026	1.92	1.02-3.63	0.043
Naïve+relapsers vs. experienced	1.11	0.68-1.81	0.663	0.41	0.24-0.69	0.001
Genotype (1b vs. 1a)	1.38	0.77-2.48	0.279	0.77	0.41-1.45	0.430
Diabetes Mellitus (no vs. yes)	1.32	0.65-2.67	0.434	1.52	0.71-3.22	0.279
Age (≥ 50 vs. < 50 years)	0.89	0.55-1.46	0.662	1.12	0.64-1.96	0.686
Fibrosis (≥F3 vs. <F3)	0.93	0.52-1.67	0.82	1.55	0.77-3.12	0.223