

In situ immunopathological changes in cutaneous leishmaniasis due to *Leishmania donovani*

Nuwani H Manamperi^a, Steve Oghumu^b, Nishantha Pathirana^c, Vipula C de Silva^d,
Wimaladharma Abeyewickreme^a, Abhay R Satoskar^e, Nadira D Karunaweera^{f*}

^a Department of Parasitology, Faculty of Medicine, University of Kelaniya, P.O. Box 06,
Thalagolla Road, Ragama, Sri Lanka

^b Department of Environmental Health Sciences, College of Public Health, Ohio State
University, Columbus, OH, USA

^c Department of Dermatology, Army Hospital, Colombo-05, Sri Lanka

^d Department of Pathology, Faculty of Medicine, University of Colombo, P. O. Box. 271, Kinsey
Road, Colombo-08, Sri Lanka

^e Department of Pathology, Wexner Medical Center, Ohio State University, Columbus, OH, USA

^f Department of Parasitology, Faculty of Medicine, University of Colombo, P. O. Box. 271,
Kinsey Road, Colombo-08, Sri Lanka

*Correspondence: Prof. Nadira D. Karunaweera, Department of Parasitology, Faculty of
Medicine, University of Colombo, P. O. Box. 271, Kinsey Road, Colombo-08, Sri Lanka. Tel.:
+94 11 2699284, Fax: +94 11 2699284, e-mail address: nadira@parasit.cmb.ac.lk


Table S1. Study and control group characteristics

	Study group (n = 58)	Control group (n = 25)	P value
Male: Female	37: 21	17: 8	0.72 ^a
Median age ^b (IQ range)	31.5 (25.7-43)	39.0 (26.5-50)	0.19 ^c

^a Pearson's Chi-Squared test; ^b age in years; ^c Mann-Whitney U test

Abbreviations: IQ, interquartile.

Figure S1. Epidermal changes seen in histopathology



Histological changes seen in the epidermis of H & E stained sections of cutaneous lesions caused by *L. donovani* in Sri Lanka. Picture A demonstrate hyperkeratosis (single arrow) and irregular acanthosis (double arrows) and picture B shows parakeratosis (single arrow) and follicular plugging (double arrows).