

**PHYSICAL ACTIVITY BUT NOT SEDENTARY ACTIVITY IS REDUCED IN PRIMARY
SJÖGREN'S SYNDROME**

Rheumatology International

Wan-Fai Ng¹, Ariana Miller², Simon J Bowman³, Elizabeth J Price⁴, George D Kitas⁵, Colin Pease⁶, Paul Emery⁶, Peter Lanyon⁷, John Hunter⁸, Monica Gupta⁸, Ian Giles⁹, David Isenberg⁹, John McLaren¹⁰, Marian Regan¹¹, Annie Cooper^{12,13}, Steven A Young-Min¹³, Neil McHugh¹⁴, Saravanan Vadivelu¹⁵, Robert J Moots¹⁶, David Coady¹⁷, Kirsten MacKay¹⁸, Bhaskar Dasgupta¹⁹, Nurhan Sutcliffe²⁰, Michele Bombardieri²⁰, Costantino Pitzalis²⁰, Bridget Griffiths²¹, Sheryl Mitchell²¹, Samira Tatiyama Miyamoto²², Michael Trenell² on behalf of the UK primary Sjögren's syndrome registry²³

¹Musculoskeletal Research Group, ²MoveLab, Physical Activity & Exercise Research, Institute of Cellular Medicine and NIHR Biomedical Research Centre for Ageing & Chronic Disease, Newcastle University, Newcastle upon Tyne, UK

³University Hospital Birmingham, Birmingham, UK

⁴Great Western Hospitals NHS Foundation Trust, Swindon, UK.

⁵Department of Rheumatology, Dudley Group of Hospitals NHS Trust, Dudley, UK.

⁶Institute of Rheumatic and Musculoskeletal Medicine, University of Leeds & NIHR Leeds Musculoskeletal Biomedical Research Unit, Leeds Teaching Hospitals Trust, Leeds, UK.

⁷Nottingham University Hospital, Nottingham, UK

⁸Gartnavel General Hospital, Glasgow, UK.

⁹University College London Hospitals NHS Foundation Trust, London, UK.

¹⁰NHS Fife, Whyteman's Brae Hospital, Kirkcaldy, UK

¹¹Royal Derby Hospital, Derby, UK.

¹²Royal Hampshire County Hospital, Winchester, UK.

¹³Portsmouth Hospitals NHS Trust, Portsmouth, UK.

¹⁴Royal National Hospital for Rheumatic Diseases, Bath, UK

¹⁵Queen Elizabeth Hospital, Gateshead, UK.

¹⁶Aintree University Hospitals, Liverpool, UK

¹⁷Royal Sunderland Hospital, Sunderland, UK

¹⁸Torbay Hospital, Torquay, UK

¹⁹Southend University Hospital, Westcliff-on-Sea, UK.

²⁰Barts and the London School of Medicine and Dentistry, UK

²¹Newcastle upon Tyne Hospitals NHS Foundation Trust, Newcastle upon Tyne, UK.

²² Universidade Federal do Espirito Santo, Vitoria, Universidade Federal de São Paulo, São Paulo, BR.

CAPES Foundation scholar – Proc. n. BEX 8831/14-9

²³Denotes corporate authorship. See appendix 1 for the full list of members.

Corresponding author: Wan-Fai Ng.

e-mail: Wan-fai.Ng@ncl.ac.uk

Supplementary Table S3. Gender characteristics of the primary Sjögren's syndrome (PSS) cohort and matched healthy control

	PSS cohort			Healthy control		
	Female	Male	p	Female	Male	p
Sample size	254	19		254	19	
Age (years)	58 (47-65)	53 (46-63)	0.372	58 (47-65)	54 (46-60)	0.404
Body mass index (kg/m ²)	25 (23-28)	27 (24-29)	0.532	25 (23-27)	26 (23-28)	0.298
Sitting time (min)	240 (120-398)	300 (240-360)	0.798	343 (223-429)	394 (266-480)	0.976
Vigorous PA (MET*min/wk)	0 (0-480)	0 (0-1920)	0.022	480 (0-1920)	960 (0-2200)	0.380
Moderate PA (MET*min/wk)	0 (0-420)	160 (0-1200)	0.048	1560 (540-3900)	1740 (960-2880)	0.391
Walking (MET*min/wk)	792 (396-2079)	990 (495-2376)	0.999	990 (462-3069)	792 (198-2079)	0.459
Total PA score (MET*min/wk)	1140 (590-3012)	3150 (1224-6750)	0.003	3690 (1732-9739)	3990 (1700-6405)	0.846

All values are presented as medians (interquartile ranges)

kg/m²: kilogram-meter squared; min: minutes; MET: metabolic equivalent of task; min/wk: minutes per week