

Supporting Information for: Direct Evidence of Lack of Colocalisation of Fluorescently Labelled Gold Labels Used in Correlative Light Electron Microscopy

Benjamin T. Miles^{1,+}, Alexander B. Greenwood^{1,+}, David Benito-Alifonso², Hugh Tanner^{3,5}, M. Carmen Galan², Paul Verkade^{4,5}, and Henkjan Gersen^{1,3,*}

¹Nanophotonics and Nanophysics Group, H.H. Wills Physics Laboratory, University of Bristol, Bristol, BS8 1TL, UK

²School of Chemistry, University of Bristol, Cantock's Close, Bristol, BS8 1TS, UK

³Bristol Centre for Functional Nanomaterials, H.H. Wills Physics Laboratory, University of Bristol, Bristol, BS8 1TL, UK

⁴Wolfson Bioimaging Facility, University of Bristol, Bristol, BS8 1TD, UK

⁵School of Biochemistry, University of Bristol, Bristol, BS8 1TD, UK

*h.gersen@bristol.ac.uk

Diffusion ordered NMR spectroscopy (DOSY) measurements were performed using a Varian VNMR500 spectrometer with a direct probe at 298 K. A pulsed gradient duration of 2 ms incremented from 3.9 to 45.5 G cm⁻¹ in 15 steps and a diffusion time of 50 ms. DOSY measurements of a sample of NP-A-633 showed the presence of a set of signals corresponding to Streptavidin-Alexa633 with diffusion coefficients between 0.3-0.5x10⁻¹⁰ ms⁻². Peaks at higher diffusion coefficients corresponding to unbound citrate (3.2x10⁻¹⁰ ms⁻²), used to stabilize gold nanoparticles, were also observed.




Figure 1. ^1H NMR of Streptavidin-Alexa633




Figure 2. DOSY of NP-A-633