

Supplement

Supplemental Table 1. Neuroticism group demographic characteristics.

	High Neuroticism Patients (N = 108)	Normal Neuroticism Patients (N = 48)	High Neuroticism Controls (N = 17)	Normal Neuroticism Controls (N = 108)	p-Value
Mean Age	32.17 ± 1.21	32.33 ± 1.69	26.12 ± 1.79	31.57 ± 1.14	0.43
Sex					0.36
Male	75	30	10	64	
Female	33	18	7	44	
Race					0.12
White	73	22	12	74	
Black/African American	28	24	4	30	
Other	7	2	1	4	
Mean Parental Education	14.04 ± 0.34	13.67 ± 0.47	15.12 ± 0.50	14.44 ± 0.20	0.04*
Mean Age at Onset of Illness	20.55 ± 0.55	22.18 ± 0.84	--	--	0.11
Mean Duration of Illness	10.26 ± 1.49	10.49 ± 2.02	--	--	0.92
Mean Chlorpromazine Equivalent	458.02 ± 31.50	491.92 ± 49.16	--	--	0.57
Mean PANSS Score					
Positive Subscale	18.46 ± 0.68	18.35 ± 0.88	--	--	0.92
Negative Subscale	15.24 ± 0.68	18.24 ± 1.23	--	--	0.04*
General Subscale	32.90 ± 0.81	30.41 ± 1.15	--	--	0.08

* significant at $\alpha=0.05$ level