

Supplementary Table 1. Estimated Caloric Intake and Dosage

		<i>Control</i>	<i>Bisphenol A</i>						<i>17α-Ethinyl Estradiol</i>		
		<i>0.0 ppm</i>	<i>0.03 ppm</i>	<i>0.3 ppm</i>	<i>3.0 ppm</i>	<i>30 ppm</i>	<i>300 ppm</i>	<i>0.0001 ppm</i>	<i>0.001 ppm</i>	<i>0.01 ppm</i>	
Caloric Intake (kcal/day)	<i>Gestation</i>	18.49 ± 5.01	17.57 ± 5.88	16.86 ± 3.96	18.54 ± 5.03	16.20 ± 5.19	17.08 ± 4.29	19.03 ± 4.19	18.80 ± 5.70	17.03 ± 4.74^a	
	<i>F1 F</i>	12.87 ± 1.30	12.44 ± 1.27	12.38 ± 1.33	12.63 ± 1.03	13.16 ± 0.95	12.60 ± 1.27	13.63 ± 1.22	13.46 ± 1.21	12.91 ± 1.12	
	<i>F1 M</i>	14.67 ± 1.42	14.04 ± 1.51	14.42 ± 1.41	15.07 ± 1.47	16.20 ± 1.64^a	14.67 ± 1.46	15.79 ± 1.99^a	14.46 ± 1.31	14.22 ± 0.99	
Est. Dose (mg/kg/day)	<i>Gestation</i>	0.00	0.005 ± 0.002	0.051 ± 0.026	0.53 ± 0.20	5.62 ± 3.49	48.3 ± 10.6	0.00002 ± 0.00001	0.00023 ± 0.00019	0.0016 ± 0.00041	
	<i>F1 F</i>	0.00	0.005 ± 0.002	0.053 ± 0.024	0.56 ± 0.25	5.52 ± 2.67	42.3 ± 10.5	0.00002 ± 0.00001	0.00019 ± 0.00007	0.0018 ± 0.00067	
	<i>F1 M</i>	0.00	0.004 ± 0.002	0.044 ± 0.017	0.43 ± 0.14	4.49 ± 1.41	40.0 ± 14.2	0.00002 ± 0.00001	0.00013 ± 0.00005	0.0014 ± 0.00052	
Exp. Dose		0.00	0.006	0.06	0.6	6	60	0.00002	0.0002	0.002	

For each exposure group, values represent mean values ± SD (n≥14 litters; Range 14-20) based on estimates of consumption of chow per animal consuming chow with a caloric density 3.63 kcal/per g. ^a Significantly different from Control (p < 0.05). Est. Dose - Estimated Dose; Exp. Dose - Expected dose based on consumption of 15% of body weight (mg/kg/day).