

Supplemental Material

Acute effect of ozone exposure on daily mortality in seven cities of Jiangsu Province, China: No clear evidence for threshold

Kai Chen, Lian Zhou, Xiaodong Chen, Jun Bi^{*}, and Patrick L. Kinney^{*}

*** Corresponding authors.**

Email address: (1) jbi@nju.edu.cn (J. Bi); (2) plk3@cumc.columbia.edu (P.L. Kinney).

Table of Contents.

Table S1. Spearman correlation coefficients between air pollutant concentrations, temperature, and relative humidity (RH) by season in seven Jiangsu cities during 2013-2014.

Fig. S1. Locations of seven study cities, air quality monitoring sites (red points), and weather stations (blue points) in Jiangsu Province, China.

Fig. S2. Monthly variations of the daily maximum 8-h average (MDA8) ozone concentrations in seven Jiangsu cities during January 18, 2013 to December 31, 2014. The lower and upper “hinges” are the 25th and 75th percentiles. The middle hinge is the median value. The upper whisker extends from the upper hinge to the highest value that is within 1.5× IQR of the upper hinge, while the lower whisker extends from the lower hinge to the lowest value that is within 1.5× IQR of the lower hinge. IQR is the inter-quantile range (75th percentile – 25th percentile). Points correspond to data beyond the end of the whiskers, i.e., outliers.

Fig. S3. Number of days with MDA8 ozone concentration exceeding the Chinese Ambient Air Quality Standard (CAAQS) level 2 standard (160 µg/m³) in seven Jiangsu cities during January 18, 2013 to December 31, 2014.

Fig. S4. Per cent increase (posterior mean and 95% posterior intervals) of daily total non-accidental mortality associated with a 10 µg/m³ increase of ozone concentration at varying lag days in seven Jiangsu cities, 2013-2014.

Fig. S5. Concentration-response curve for ozone (lag01) and daily total mortality based on natural cubic spline models with 3df in seven Jiangsu cities, 2013-2014. Linearity p-value represents the significance of spline models departure from linear models.

Table S1. Spearman correlation coefficients between air pollutant concentrations, temperature, and relative humidity (RH) by season in seven Jiangsu cities during 2013-2014.

	PM _{2.5}	PM ₁₀	NO ₂	SO ₂	Temperature	RH
<i>Full year</i>						
O ₃	-0.053	-0.003	-0.136	-0.037	0.396	-0.119
PM _{2.5}		0.893	0.639	0.617	-0.321	-0.166
PM ₁₀			0.637	0.682	-0.272	-0.371
NO ₂				0.627	-0.293	-0.32
SO ₂					-0.366	-0.552
Temperature						0.143
<i>Warm season (Apr.-Sep.)</i>						
O ₃	0.306	0.37	0.195	0.338	0.085	-0.426
PM _{2.5}		0.873	0.563	0.493	-0.103	-0.202
PM ₁₀			0.557	0.607	-0.07	-0.433
NO ₂				0.585	-0.133	-0.34
SO ₂					-0.007	-0.616
Temperature						-0.122
<i>Cold season(Oct.-Mar.)</i>						
O ₃	-0.108	-0.105	-0.179	-0.134	0.327	0.073
PM _{2.5}		0.897	0.606	0.632	-0.092	-0.06
PM ₁₀			0.623	0.676	-0.046	-0.259
NO ₂				0.565	0.079	-0.258
SO ₂					-0.274	-0.474
Temperature						0.134




Fig. S1. Locations of seven study cities, air quality monitoring sites (red points), and weather stations (blue points) in Jiangsu Province, China.




Fig. S2. Monthly variations of the daily maximum 8-h average (MDA8) ozone concentrations in seven Jiangsu cities during January 18, 2013 to December 31, 2014. The lower and upper “hinges” are the 25th and 75th percentiles. The middle hinge is the median value. The upper whisker extends from the upper hinge to the highest value that is within 1.5× IQR of the upper hinge, while the lower whisker extends from the lower hinge to the lowest value that is within 1.5× IQR of the lower hinge. IQR is the inter-quantile range (75th percentile – 25th percentile). Points correspond to data beyond the end of the whiskers, i.e., outliers.




Fig. S3. Number of days with MDA8 ozone concentration exceeding the Chinese Ambient Air Quality Standard (CAAQS) level 2 standard ($160 \mu\text{g}/\text{m}^3$) in seven Jiangsu cities during January 18, 2013 to December 31, 2014.




Fig. S4. Per cent increase (posterior mean and 95% posterior intervals) of daily total non-accidental mortality associated with a $10 \mu\text{g}/\text{m}^3$ increase of ozone concentration at varying lag days in seven Jiangsu cities, 2013-2014.




Fig. S5. Concentration-response curve for ozone (lag01) and daily total mortality based on natural cubic spline models with 3df in seven Jiangsu cities, 2013-2014. Linearity p-value represents the significance of spline models departure from linear models.