

Supplemental informations for

Intestinal microbiome in children with severe and complicated acute viral

Gastroenteritis

Shih-Yen Chen¹, Chi-Neu Tsai², Yun-Shien Lee³, Chun-Yuan Lin⁴, Kuan-Yeh

Huang⁵, Hsun-Ching Chao¹, Ming-Wei Lai¹ & Cheng-Hsun Chiu^{2,5,*}

¹Division of Pediatric Gastroenterology, Chang Gung Children's Hospital, Chang Gung University College of Medicine, Taoyuan, Taiwan.


²Graduate Institute of Clinical Medical Sciences, Chang Gung University College of Medicine, Taoyuan, Taiwan

³Department of Biotechnology, Min-Chuan University, Taoyuan, Taiwan

⁴Department of Computer Science and Information Engineering, Chang Gung University College of Engineering, Taoyuan, Taiwan


⁵Molecular Infectious Disease Research Center, Chang Gung Memorial Hospital, Chang Gung University College of Medicine, Taoyuan, Taiwan.

sFig. 1A




Intestinal microbiota in a 37-day-old neonate with noroviral AGE and necrotizing enterocolitis.

sFig. 1B



Intestinal microbiota in a 9-month-old male children with rotavirus AGE and convulsion.

sFig. 1C



Intestinal microbiota in an 11-month-old male children with noroviral AGE and acute renal failure.