

# **Establishment and characterization of an orthotopic patient-derived Group**

## **3 medulloblastoma model for preclinical drug evaluation**

Emma Sandén<sup>1,a</sup>, Cecilia Dyberg<sup>2,a</sup>, Cecilia Krona<sup>3</sup>, Gabriel Gallo-Oller<sup>2</sup>, Thale Kristin Olsen<sup>2</sup>, Julio Enríquez Pérez<sup>1</sup>, Malin Wickström<sup>2</sup>, Atosa Estekizadeh<sup>4</sup>, Marcel Kool<sup>5</sup>, Edward Visse<sup>1</sup>, Tomas J. Ekström<sup>4</sup>, Peter Siesjö<sup>1,6</sup>, John Inge Johnsen<sup>2,b</sup>, Anna Darabi<sup>1,b\*</sup>

\* corresponding author

<sup>a</sup>shared first authorship

<sup>b</sup>shared senior authorship

<sup>1</sup>Lund University, Faculty of Medicine, Department of Clinical Sciences Lund, Neurosurgery, Lund, Sweden

<sup>2</sup>Karolinska Institutet, Department of Women's and Children's Health, Childhood Cancer Research Unit, Stockholm, Sweden

<sup>3</sup>Uppsala University, Department of Immunology, Genetics and Pathology, Uppsala, Sweden

<sup>4</sup>Karolinska University Hospital, Solna, Center for Molecular Medicine, and Karolinska Institutet, Department of Clinical Neuroscience, Stockholm, Sweden

<sup>5</sup>German Cancer Research Center DKFZ, Division of Pediatric Neurooncology, Heidelberg, Germany

<sup>6</sup>Lund University, Skane University Hospital, Department of Clinical Sciences Lund, Neurosurgery, Lund, Sweden



