Translation, but not transfection limits clinically relevant, exogenous mRNA based induction of

alpha-4 integrin expression on human mesenchymal stem cells

Adam Nowakowski¹, Anna Andrzejewska¹, Johannes Boltze^{2,3,4,5}, Franziska Nitzsche^{2,3}, Li-li Cui6, Jukka

Jolkkonen^{6,7}, Piotr Walczak^{7,8,9}, Barbara Lukomska¹, Miroslaw Janowski^{1,8,10}

1 NeuroRepair Department, Mossakowski Medical Research Centre, Polish Academy of Sciences, Warsaw,

Poland, 2 Fraunhofer-Institute for Cell Therapy, and Immunology, Department of Cell Therapy,

University of Leipzig, Leipzig, Germany, 3 Translational Centre for Regenerative Medicine, University of

Leipzig, Leipzig, Germany, 4 Fraunhofer Research Institution of Marine Biotechnology, Department of

Translational Medicine, University of Lubeck, Lubeck, Germany, 5 Cell Technology, Lübeck, German;

Institute for Medical and Marine Biotechnology, University of Lübeck, Lübeck, Germany, 6 Stroke Recovery

Laboratory, Institute of Clinical Medicine - Neurology, University of Eastern Finland, Kuopio, Finland,

7 Neurocenter, Neurology, University Hospital of Kuopio, Kuopio, Finland, 8 Russell H. Morgan Dept.

of Radiology and Radiological Science, Division of MR Research, 9 Institute for Cell Engineering, The

Johns Hopkins University School of Medicine, Baltimore, USA, 10 Department of Radiology, Faculty

of Medical Sciences, University of Warmia and Mazury, Olsztyn, Poland

*Corresponding author: mjanows1@jhmi.edu

running head: mRNA-based approach to alpha-4 integrin

keywords: mRNA, ARCA, transfection, stroke, mesenchymal stem cells

Nowakowski et al.

- **Supplementary figure 1. Overview of the experiments (graphical abstract)**
- 2 Supplementary figure 2. pITGA4-SP72 plasmid template DNA preparation with subsequent
- 3 verification of ITGA4-mRNA production in vitro
- 4 E.coli were transfected with pITGA4-SP72 and obtained selected clones were digested with
- 5 XbaI/XhoI to confirm proper subcloning from pITGA4 to pSP72 (A). The cloning site was verified by
- 6 sequencing (B). Selected clone pITGA4-SP72 was employed for ITGA4-mRNA production in vitro
- 7 (C).
- 8 Supplementary figure 3. ITGA4 protein synthesis after ITGA4-mRNA transfection
- 9 Immunocytochemical assessment of ITGA4 protein presence after ITGA4-mRNA bearing natural 7-
- methylguanosine cap transfection using Lipofectamine®2000.

11









