

Supplementary Information for:

Chemokine profiles of interstitial pneumonia in patients with dermatomyositis: a case control study

Katsuhiko Oda, MD¹, Takuya Kotani, MD, PhD¹, Tohru Takeuchi, MD, PhD¹, Takaaki Ishida, MD¹, Takeshi Shoda, MD¹, Kentaro Isoda, MD, PhD¹, Shuzo Yoshida, MD, PhD¹, Yasuichiro Nishimura, PhD², and Shigeki Makino, MD, PhD¹

¹Department of Internal Medicine (I), Osaka Medical College, Takatsuki, Osaka, Japan

²Department of Mathematics, Osaka Medical College, Takatsuki, Osaka, Japan

Correspondence and reprint requests to: Takuya Kotani, MD, PhD,

Department of Internal Medicine (I), Osaka Medical College, Daigaku-Machi 2-7,
Takatsuki, Osaka 569-8686, Japan


Tel.: +81-72-683-1221; Fax: +81-72-683-1801

E-mail: in1242@osaka-med.ac.jp

This supplement contains:

Supplementary Figure S1

Supplementary Fig. S1. Comparison of the change ratio of chemokine levels initially and at 2 weeks after treatment.



CCL: C-C motif chemokine ligand; CXCL: C-X-C motif chemokine ligand. The P value was estimated by the Mann-Whitney U-test. * $P < 0.05$.