

Table S2 Stomatal conductance (g_s , H_2O mol $\text{m}^{-2} \text{s}^{-1}$) in the WW, TD, and SD vines during the acclimation period (Fig. 2b). Different letters for each row (DOE = Days of Experiment) denotes significant differences ($P<0.05$) between treatments as tested by Tukey HSD

DOE	WW	TD	SD
0	0.349 a	0.324 a	0.362 a
2	0.249 a	0.173 a	0.219 a
5	0.216 a	0.188 a	0.217 a
7	0.226 a	0.156 b	0.182 ab
9	0.281 a	0.215 a	0.228 a
11	0.374 a	0.294 ab	0.215 b
13	0.378 a	0.137 b	0.080 c
14	0.413 a	0.250 b	0.151 c
16	0.273 a	0.198 b	0.057 c
18	0.425 a	0.354 b	0.180 c
19	0.375 a	0.341 a	0.101 b
23	0.293 a	0.305 a	0.053 b
28	0.563 a	0.089 b	0.120 b
29	0.412 a	0.056 b	0.079 b
30	0.455 a	0.159 b	0.187 b
33	0.457 a	0.319 a	0.176 b
34	0.422 a	0.335 b	0.080 c
37	0.272 a	0.250 a	0.033 b
39	0.278 a	0.034 b	0.028 b
40	0.296 a	0.081 b	0.101 b
42	0.388 a	0.287 b	0.225 b
46	0.309 a	0.258 a	0.271 a