Table S1. Humanized mice used in Figures 1 and 2.

Mouse no.	Recipient mouse		Transplanted hHSCs§		ADORECOLI bonlotuno		Inoculated age
	Lot no.*	Sex†	Donor lot‡	No. of cells	APOBEC3H haplotype		(weeks)
1	235	F	Α	150,000	III / <b>V</b>	Stable	18
2	235	M	Α	150,000	III / <b>V</b>	Stable	18
3	238	F	В	120,000	1 / <b>II</b>	Stable	13
4	238	F	В	120,000	1 / <b>II</b>	Stable	13
5	238	M	В	120,000	1 / <b>II</b>	Stable	13
6	243	M	В	230,000	1 / <b>II</b>	Stable	14
7	243	M	В	230,000	1 / <b>II</b>	Stable	14
8	243	M	В	230,000	1 / <b>II</b>	Stable	14
9	234	M	С	135,000	1/1	Intermediate	18
10	253	M	С	160,000	1/1	Intermediate	15
11	253	M	С	160,000	1/1	Intermediate	15
12	253	M	С	160,000	1/1	Intermediate	15
13	241	F	D	140,000	I / VI	Intermediate	14
14	250	M	Е	100,000	1/1	Intermediate	14

<sup>\*</sup> Seven lots of newborn NOG mice were used for the recipient.

<sup>†</sup> F, female; M, male.

<sup>‡</sup> NOG-hCD34 mice were reconstructed with one of 5 donors.

<sup>§</sup> hHSCs, human CD34<sup>+</sup> hematopoietic stem cells.