

Supplementary Table S1. Sample Characteristics for Mothers of Girls and Girls at 9 Years of Age with PFASs and DXA

Variable	N	Median (IQR)
PFOA (ng/mL)	359	3.7 (2.9–4.8)
PFOS (ng/mL)	359	19.8 (15.0–25.3)
PFHxS (ng/mL)	359	1.6 (1.3–2.2)
PFNA (ng/mL)	358	0.5 (0.4–0.7)
Gestational age at blood collection (weeks)	359	15 (10–28)
Maternal age at collection (years)	359	28.8 (25.6–32.1)
Girls' DXA-derived total body fat at age 9 (%)	359	27.5 (21.7–34.6)
Girls' DXA-derived trunk body fat at age 9 (%)	359	39.6 (36.5–43.7)
Girls' BMI at age 9 (kg/m ²)	351	18.1 (16.3–20.6)
Girls' WC at age 9 (cm)	359	62.7 (58.6–68.8)
Girls dietary intake at age 7	340	
Energy (kcal)		1649 (1426–1857)
Fat (% energy)		36.7 (33.5–39.7)
Carbohydrate (% energy)		53.5 (50.0–57.1)
Protein (% energy)		13.0 (11.5–14.5)

DXA, dual energy X-ray absorptiometry; IQR, interquartile range; PFHxS, perfluorohexane sulfonate; PFNA, perfluorononanoate; PFOA, perfluorooctanoate; PFOS, perfluorooctane sulfonate; WC, waist circumference.