

Supplementary Table S2. Regression Coefficients (β) for the Relationship between Perfluoroalkyl Substances and Measures of Body Fatness (Unadjusted) Stratified by Educational Status

Analyte	DXA-total body fat (%)		
	β	95% CI	p
PFOS (ng/mL)			
Overall ^a	-0.03	-0.11 to 0.05	0.47
Low education	0.03	-0.11 to 0.17	0.71
Medium	0.12	-0.04 to 0.27	0.13
High	-0.18	-0.30 to -0.06	0.002
PFOA (ng/mL)			
Overall ^a	-0.17	-0.60 to 0.26	0.44
Low education	-0.22	-1.13 to 0.68	0.63
Medium	1.42	0.38 to 2.45	0.007
High	-0.51	-1.05 to 0.02	0.06
PFHxS (ng/mL)			
Overall ^a	-0.03	-0.16 to 0.10	0.65
Low education	-0.07	-0.34 to 0.20	0.59
Medium	0.19	-0.12 to 0.49	0.23
High	-0.09	-0.26 to 0.08	0.29
PFNA (ng/mL)			
Overall ^a	2.12	-0.89 to 5.12	0.17
Low education	16.32	4.31 to 28.33	0.008
Medium	2.93	-2.10 to 7.97	0.25
High	-0.06	-3.68 to 3.57	0.97

^aOverall N = 359; N = 61, N = 115, and N = 170 for the low, medium, and high educated groups, respectively, with N = 13 missing educational status.