

Parameters common to all simulations

Parameter	Value	Units	Reference
Cell volume	10^{-18}	m^{-3}	Bremer and Dennis (1996); Elf et al. (2003)
Cell doubling time	30	min	This work
Concentration of tRNAs	4.08×10^5	molecules/cell	Bremer and Dennis (2008)
Rate of peptide-bond formation and translocation	≈ 22 (1/0.045)	s^{-1}	Bremer and Dennis (2008)
tRNA accommodation rate at normal sites (not stall sites)	80–1650	s^{-1}	Inferred (Methods)
Ribosome footprint size on mRNAs	10	codons	This work
Number of distinct mRNA species	2	No units	This work
Ribosome-tRNA association rate constant	2×10^7	$\text{M}^{-1} \text{s}^{-1}$	Bilgin and Ehrenberg (1994); Pavlov and Ehrenberg (1996)
tRNA aminoacylation rate constant	2×10^{10}	$\text{M}^{-1} \text{s}^{-1}$	This work
Threshold tRNA accommodation rate for selective abortive termination	22	s^{-1}	This work
Abortive termination rate constant (SAT and CSAT models)	1	s^{-1}	This work
Abortive termination rate constant (TJ model)	0	s^{-1}	This work
Translation initiation rate	0.3	s^{-1}	This work

References

- Bilgin N, Ehrenberg M. Mutations in 23 S Ribosomal RNA Perturb Transfer RNA Selection and Can Lead to Streptomycin Dependence. *Journal of Molecular Biology*. 1994 jan; 235(3):813–824.
- Bremer H, Dennis P. Modulation of Chemical Composition and Other Parameters of the Cell at Different Exponential Growth Rates. *EcoSal Plus*. 2008; doi: [10.1128/ecosal.5.2.3](https://doi.org/10.1128/ecosal.5.2.3).
- Bremer H, Dennis P. Modulation of Chemical Composition and Other Parameters of the Cell by Growth Rate. In: *Escherichia Coli and Salmonella* Washington, DC: ASM Press; 1996.p. 1553–1569.
- Elf J, Nilsson D, Tenson T, Ehrenberg M. Selective Charging of tRNA Isoacceptors Explains Patterns of Codon Usage. *Science*. 2003 jun; 300(5626):1718–1722.
- Pavlov MY, Ehrenberg M. Rate of Translation of Natural mRNAs in an Optimized in Vitro System. *Archives of Biochemistry and Biophysics*. 1996 apr; 328(1):9–16.