


THE LANCET

Supplementary appendix


This appendix formed part of the original submission and has been peer reviewed. We post it as supplied by the authors.

Supplement to: WOMAN Trial Collaborators. Effect of early tranexamic acid administration on mortality, hysterectomy, and other morbidities in women with post-partum haemorrhage (WOMAN): an international, randomised, double-blind, placebo-controlled trial. *Lancet* 2017; published online April 26. [http://dx.doi.org/10.1016/S0140-6736\(17\)30638-4](http://dx.doi.org/10.1016/S0140-6736(17)30638-4).

Supplementary file 1a. Cause of death by hours from giving birth.



Supplementary file 1b. Cause of hysterectomy by hours from giving birth.



Adverse Events

Event	TXA	Placebo
Postpartum bleeding (secondary)	6	4
Transfusion reaction	3	1
Anaemia, postpartum	2	2
Retention urine	2	1
Urinary tract infection	2	2
Wound infection	2	1
Abdominal pain	1	0
Allergic rash	1	0
Calf pain	1	0
Endometritis postpartum	1	1
Haematoma	1	0
Headache	1	0
Hypoglycaemic episode	1	0
Incontinence of urine	1	0
Pulmonary embolism	1	1
Thrombotic thrombocytopenic purpura	1	0
Vaginal infection	1	0
Vesicovaginal fistula	1	0
Vomiting	1	0
Allergic reaction	0	1
Anxiety	0	1
Cardiac arrest	0	1
Disruption of caesarean wound	0	1
Dysfunctional uterine bleeding	0	1
Eclampsia	0	1
Haematuria	0	1
Jaundice	0	1
Malaria	0	1
Papilloedema	0	1
Perineal infection	0	1
Retained products of conception	0	1
Swelling	0	1
Urticarial rash	0	1