

**Characterisation and antimicrobial activity of biosurfactant extracts
produced by *Bacillus amyloliquefaciens* and *Pseudomonas
aeruginosa* isolated from a wastewater treatment plant**

**Thando Ndlovu¹, Marina Rautenbach², Johann Arnold Vosloo², Sehaam Khan³ and
Wesaal Khan^{1*}**

¹Department of Microbiology, Faculty of Science, Stellenbosch University, Private Bag X1,
Stellenbosch, 7602, South Africa.

²BIOPEP Peptide Group, Department of Biochemistry, Faculty of Science, Stellenbosch
University, Private Bag X1, Stellenbosch, 7602, South Africa.

³Faculty of Health and Applied Sciences, Namibia University of Science and Technology, 13
Storch Street, Private Bag 13388, Windhoek, Namibia.

*Corresponding Author: Wesaal Khan, Email: wesaal@sun.ac.za, Tel: +27 21 808 5804,
Fax: +27 21 808 5846




Fig. S1 Comparison of the different culture extracts showing the relative contribution of each of the surfactin groups in the biosurfactant extracts. The contribution was calculated from UPLC profiles, with the assumption that all the surfactin species has similar ion responses. SARCC 697 LC = glycerol-MSM culture extract, SARCC 697 LC-AE = 70% acetonitrile extract of SARCC 697 LC, SARCC 697 NA-TSC = NA test tube slant culture extract, SARCC 697 NA-PDC = NA petri dish culture extract

