

Supplemental Digital Content 1

Table S1. Bivariate associations among MetS components and SES variables (N = 354).

	1	2	3	4	5	6	7	8	9	10	11
1. Early-life SES	1	.07	.13*	.14**	.14**	.15**	.04	.07	.09 ^Δ	-.05	.06
2. Current SES		1	.09 ^Δ	.09 ^Δ	.05	.07	-.07	-.03	.03	-.09 ^Δ	.01
3. MetS diagnosis			1	.67**	.58**	.47**	.31**	.34**	.49**	-.21**	.43**
4. Number of MetS components				1	.78**	.69**	.45**	.48**	.53**	-.43**	.41**
5. MetS factor score					1	.86**	.62**	.69**	.73**	-.48**	.47**
6. Waist circumference						1	.35**	.49**	.48**	-.39**	.23**
7. Diastolic blood pressure							1	.78**	.35**	-.11*	.32**
8. Systolic blood pressure								1	.33**	-.19**	.29**
9. Triglycerides									1	-.24**	.34**
10. HDL cholesterol										1	-.08
11. HbA1c											1

* $p < .05$; ** $p < .01$; ^Δ $p < .10$.




Figure S1. Bars depict the percentage of participants within each group exhibiting each count of MetS components (0, 1, 2, 3 or more MetS components).