

Plasmacytoid dendritic cells appear inactive during sub-microscopic *Plasmodium falciparum* blood-stage infection, yet retain their ability to respond to TLR stimulation.


Jessica R. Loughland^{1*}, Gabriela Minigo¹, Derek S. Sarovich^{1,2}, Matt Field³, Peta E. Tipping^{1€}, Marcela Montes de Oca⁴, Kim A. Piera¹, Fiona H. Amante⁴, Bridget E. Barber^{1,6}, Matthew J. Grigg^{1,6}, Timothy William^{6,7}, Michael F. Good⁸, Denise L. Doolan^{3,4}, Christian R. Engwerda⁴, Nicholas M. Anstey^{1,5}, James S. McCarthy⁴⁺, Tonia Woodberry¹⁺.

¹Menzies School of Health Research, Darwin, Australia and Charles Darwin University, Darwin, Australia; ²Centre for Animal Health Innovation, Faculty of Science, Health, Education and Engineering, University of the Sunshine Coast, Sippy Downs, Queensland, Australia; ³Australian Institute of Tropical Health and Medicine, James Cook University, Cairns, Australia; ⁴QIMR Berghofer Medical Research Institute, Brisbane, Australia; ⁵Royal Darwin Hospital, Darwin, Australia; ⁶Infectious Diseases Unit, Queen Elizabeth Hospital, Kota Kinabalu, Sabah, Malaysia; ⁷Sabah Department of Health, Kota Kinabalu, Sabah, Malaysia; ⁸Griffith University, Gold Coast, Australia.

* jessica.loughland@menzies.edu.au

+ These authors contributed equally to this work.

€ Present Address: Royal Darwin Hospital, Darwin, Australia.



Supplementary Figure 1 Summary of differential gene expression. Longitudinal pDC gene expression changes in five participants, between day 0 and peak parasitemia (day 7-8). Represented genes had an FDR<0.07, each symbol represents one participant.