

**Pluronic F127 and D- α -Tocopheryl Polyethylene Glycol Succinate (TPGS)
Mixed Micelles for Targeting Drug Delivery
across The Blood Brain Barrier**

Xin Meng^{1*}, Jiansheng Liu^{2*}, Xiangrong Yu^{3*}, Jiajia Li¹, Xiaotong Lu¹, Teng Shen^{1,4}

¹Department of Pharmaceutics, Key Laboratory of Smart Drug Delivery, Ministry of Education, School of Pharmacy, Fudan University, 826 Zhangheng Road, Shanghai 201203, China

²Department of Neurology, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine, 639 Zhizaoju Road, Shanghai 200011, China

³Department of Radiology, Zhuhai People's Hospital, Zhuhai Hospital of Jinan University, 79 Kangning Road, Zhuhai 519000, China

⁴The Institutes of Integrative Medicine of Fudan University, 12 Wulumuqi Middle Road, Shanghai 200040, China

*These authors contributed equally to this work.

Corresponding author: Teng Shen, Department of Pharmaceutics, Key Laboratory of Smart Drug Delivery, Ministry of Education, School of Pharmacy, Fudan University; The Institutes of Integrative Medicine of Fudan University.

Address: 826 Zhangheng Road, Shanghai 201203, China; 12 Wulumuqi Middle Road, Shanghai 200040, China.

Tel.: +86-021-51980081; fax: +86-021-51980081.

E-mail: shenteng@fudan.edu.cn.

Supplementary data.




Figure S1. The stability of DiR-loaded F127/TPGS mixed micelles with different ratios incubated at 37 °C. Each point represents mean \pm SD (n = 3).