

**Extrasynaptic NMDA receptor dependent long-term potentiation of
hippocampal CA1 pyramidal neurons**

Qian Yang^{1#}, Geng Zhu^{1#}, Dandan Liu¹, Jue-Gang Ju¹, Zhen-Hua Liao¹, Yi-Xin Xiao¹,
Yue Zhang¹, Naijian Chao¹, JieJie Wang², Weidong Li¹, Jian-Hong Luo^{2*},
Sheng-Tian Li^{1*}

¹Key laboratory for the Genetics of Developmental and Neuropsychiatric Disorders
(Ministry of Education), Bio-X Institutes, Shanghai Key Laboratory of Psychotic
Disorders, Institute of Social Cognitive and Behavioral Sciences, and Brain Science
and Technology Research Center, Shanghai Jiao Tong University, Shanghai, China.

²Department of Neurobiology, Key Laboratory of Medical, Neurobiology (Ministry of
Health of China), Collaborative Innovation Center for Brain Science, School of
Medicine, Zhejiang University, Hangzhou, Zhejiang 310058, China

* luojianhong@zju.edu.cn; Tel.: +86-571-88208002; Fax: +86-571-88208002

* lstian@sjtu.edu.cn; Tel.: +86-21-34208036; Fax: +86-21-34208035

contributed equally to this work.

Figure 6




Figure S1. Full-length blots for the indicated figures from the main text.