

Supplementary Table 2. Stratification analyses on associations between family history of stroke and incident stroke

Subgroup	HR*	95% CI*	<i>P</i> for heterogeneity	<i>P</i> for interaction
Residential area			0.145	
Rural	1.37	1.32–1.43		
Urban	1.31	1.26–1.37		
Marital status			0.012	
Married	1.36	1.32–1.40		
Widowed/separated/divorced/never married	1.22	1.13–1.32		
Education			<0.001	<0.001
Illiteracy or primary school	1.26	1.21–1.31		
Middle school and above	1.44	1.38–1.50		
Smoking status			<0.001	<0.001
Not current	1.30	1.26–1.34		
Current	1.46	1.38–1.54		
Alcohol consumption			0.005	<0.001
Not current	1.31	1.27–1.36		
Current	1.45	1.36–1.53		
Physical activity (MET h/day)			<0.001	<0.001
<18.20	1.29	1.25–1.34		
≥18.20	1.46	1.39–1.54		
BMI (kg/m ²)			0.788	
<24.0	1.34	1.28–1.40		
24.0, 27.9	1.36	1.30–1.42		
≥28.0	1.32	1.23–1.42		
Hypertension			0.035	
No	1.39	1.32–1.46		
Yes	1.30	1.26–1.35		
Diabetes			0.034	
No	1.36	1.32–1.40		
Yes	1.24	1.14–1.34		

HR, hazard ratio; CI, confidence interval; MET, metabolic equivalent of task; BMI, body mass index.

*Adjusted for age, sex, marital status, education, residential area, smoking status, alcohol consumption, physical activity, BMI, history of hypertension, history of diabetes and siblings except for the stratification factor.