

Platelet distribution width correlates with prognosis of non-small cell lung cancer

running heading: PDW and prognosis in NSCLC

Ming-ming Cui^{1#}; Na Li^{1#}; Xing Liu^{2#}, Zhi-yuan Yun¹; Ye Niu³; Yong Zhang⁴; Boning Gao⁵; Tiemin Liu^{1,6}; Rui-tao Wang^{1,*}.

¹Department of Internal Medicine, Harbin Medical University Cancer Hospital, Harbin Medical University, Harbin, Heilongjiang, 150081, China

²Department of Endocrinology, Ansteel Group Hospital, AnShan, Liaoning, 114001, China

³Department of Geriatrics, the Second Affiliated Hospital, Harbin Medical University, Harbin, Heilongjiang, 150086, China

⁴Department of Pharmacology, Harbin Medical University, Harbin, 150081, China

⁵Hamon Center for Therapeutic Oncology Research, Department of Internal Medicine Pharmacology, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA

⁶Division of Hypothalamic Research, Department of Internal Medicine, UT Southwestern Medical Center, Dallas, TX, 75390, USA

*To whom correspondence should be addressed: Rui-tao Wang. Department of Internal Medicine, Harbin Medical University Cancer Hospital, Harbin Medical University, NO.150 Haping ST, Nangang District, Harbin, 150081, China; Tel: 86-451-86298352; Fax: 86-451-86298353; Email: ruiwang@126.com.

Ming-ming Cui, Na Li, and Xing Liu contributed equally to this work.

word count: 1798; the number of figures and tables: 5

Supplementary Table 1

Correlation coefficient (r) for PDW in relation to WBC, neutrophil, platelet count, and MPV in NSCLC.

	WBC		Neutrophil		Platelet count		MPV	
	r	p	r	p	r	p	r	p
PDW	-0.137	0.024	-0.156	0.010	-0.211	<0.001	0.073	0.229