

Appendix 2. Oxycodone Utilization After Discharge and Satisfaction With Pain Regimen, Stratified by Type and Timing Of Cesarean Delivery

Outcomes	Median [IQR], or n (%)			
	Primary cesarean delivery (n=35)	Repeat cesarean delivery (n=15)	Planned ^a cesarean delivery (n=19)	Unplanned cesarean delivery (n=31)
Number of oxycodone tablets chosen,	20.0 [15.0, 25.0] ^b	20.0 [10.0, 30.0] ^c	20.0 [10.0, 23.0] ^d	20.0 [20.0, 30.0] ^e
Number of oxycodone tablets used	16.0 [8.0, 25.0]	15.0 [3.0, 30.0]	11.0 [3.0, 23.0]	16.0 [8.0, 31.0]
Number of oxycodone tablets remaining	4.0 [0.0,11.0]	0.0 [0.0, 8.0]	1.0 [0.0, 9.0]	4.0 [0.0, 8.0]
Need for oxycodone refills	2 (5.7)	2 (13.3)	0 (0.0)	4 (12.9)
Satisfied with outpatient pain management	18 (51.4)	8 (53.3)	9 (47.4)	17 (54.8)
Very satisfied with outpatient pain management	13 (37.1)	6 (40.0)	8 (42.1)	11 (35.5)

^a Planned cesarean delivery refers to a scheduled cesarean delivery. Unplanned cesarean delivery refers to an intrapartum cesarean delivery.

^b For 5 patients, the number of tablets prescribed was different from the number chosen, such that the overall median [IQR] number of tablets dispensed was 23.0 [20.0, 35.0].

^c For 1 patient, the number of tablets prescribed was different from the number chosen, such that the overall median [IQR] number of tablets dispensed was 20.0 [10.0, 30.0].

^d For 4 patients, the number of tablets prescribed was different from the number chosen, such that the overall median [IQR] number of tablets dispensed was 20.0 [10.0, 25.0].

^e For 2 patients, the number of tablets prescribed was different from the number chosen, such that the overall median [IQR] number of tablets dispensed was 25.0 [20.0, 35.0].