

**Table S7. Association of Baseline Appetite with ADMA and SDMA Concentrations in HEMO Study**

Description	N (%)	ADMA, $\mu\text{M}$		SDMA, $\mu\text{M}$	
		Mean $\pm$ SD	P <sup>1</sup>	Mean $\pm$ SD	P <sup>1</sup>
<b>During the past week (7 days), how would you rate your appetite?</b>					
Very good	390 (30.6)	0.904 $\pm$ 0.232	0.01	4.273 $\pm$ 1.431	0.8
Good	477 (37.4)	0.924 $\pm$ 0.238		4.254 $\pm$ 1.400	
Fair	301 (23.6)	0.933 $\pm$ 0.227		4.225 $\pm$ 1.298	
Poor	86 (6.7)	0.955 $\pm$ 0.180		4.402 $\pm$ 1.250	
Very poor	22 (1.7)	0.971 $\pm$ 0.281		4.285 $\pm$ 1.566	
<b>During the past 4 weeks, to what extent were you bothered by lack of appetite?</b>					
Not at all	648 (51.5)	0.916 $\pm$ 0.242	0.3	4.243 $\pm$ 1.381	0.8
Somewhat	312 (24.8)	0.914 $\pm$ 0.195		4.303 $\pm$ 1.296	
Moderately	135 (10.7)	0.943 $\pm$ 0.242		4.205 $\pm$ 1.305	
Very much	100 (7.9)	0.950 $\pm$ 0.244		4.146 $\pm$ 1.291	
Extremely	47 (3.7)	0.938 $\pm$ 0.224		4.341 $\pm$ 1.603	

Abbreviation: ADMA, Asymmetric Dimethylarginine; SDMA, Symmetric Dimethylarginine; SD, standard deviation

<sup>1</sup> P-value by Wilcoxon-Mann-Whitney test