Table S2 - List of multilocus genotypes (MLGs) across bitter manioc varieties in different soil types in Manicoré, Amazonas, Brazil. For the detection of MLGs individuals with missing data were removed. N = number of individuals. Acronyms of the varieties are presented in Table 1 of the main text.

		Shared MLGs														**Unique MLGs**																					
Varieties	**N**	**11**	**4**	**25**	**5**	**1**	**8**	**20**	**30**	**17**	**34**	**10**	**19**	**26**	**35**	**2**	**3**	**6**	**7**	**9**	**12**	**13**	**14**	**15**	**16**	**18**	**21**	**22**	**23**	**24**	**27**	**28**	**29**	**31**	**32**	**33**
PB ADE	18									17								1																		
PB FP	20	19																		1																
PA ADE	30		28								1				2																					
PA FP	29		29																																	
AR ADE	20			12						5											1			1								1				
AR OX	29			28																										1						
TA ADE	30	29	1																																	
TA FP	30							25					2													1	1	1								
TA OX	30	25	1									3																	1							
MJ FP	30				2				25							1	1						1													
OR FP	30					27																1			1									1		
AU OX	29						27												1														1			
JÁ OX	29			4							20			2																1				1	1
RO OX	30				30																															
Total	**384**	**73**	**59**	**44**	**32**	**27**	**27**	**25**	**25**	**22**	**21**	**3**	**2**	**2**	**2**	**1**	**1**	**1**	**1**	**1**	**1**	**1**	**1**	**1**	**1**	**1**	**1**	**1**	**1**	**1**	**1**	**1**	**1**	**1**	**1**	**1**