

Supplementary Table 1. Response Rates to Interferon-Based Antiviral Therapy among Korean Patients with Chronic Hepatitis C Virus Infection

	GT 1		GT 2		GT 3		GT 4		GT 6	
	ITT (n=251)	PP (n=189)	ITT (n=273)	PP (n=238)	ITT (n=6)	PP (n=5)	ITT (n=2)	PP (n=2)	ITT (n=3)	PP (n=2)
RVR	94 (37.5)	78 (41.3)	143 (52.4)	133 (55.9)	2 (33.3)	2 (40.0)	1 (50.0)	1 (50.0)	0	0
EVR	192 (76.5)	160 (84.7)	126 (46.2)	113 (47.5)	1 (16.7)	1 (20.0)	1 (50.0)	1 (50.0)	3 (100.0)	2 (100.0)
pEVR	36 (14.3)	24 (12.7)	13 (4.8)	10 (4.2)	0	0	0	0	0	0
cEVR	156 (62.2)	136 (72.0)	113 (41.4)	103 (43.3)	1 (16.7)	1 (20.0)	1 (50.0)	1 (50.0)	3 (100.0)	2 (100.0)
ETR	185 (73.7)	171 (90.5)	223 (81.7)	211 (88.7)	5 (83.3)	5 (100.0)	2 (100.0)	2 (100.0)	2 (66.7)	2 (100.0)
SVR	147 (58.6)	135 (71.4)	203 (74.4)	193 (81.1)	5 (83.3)	5 (100.0)	1 (50.0)	1 (50.0)	2 (66.7)	2 (100.0)
Breakthrough	9 (3.6)	5 (2.6)	4 (1.5)	4 (1.7)	0	0	0	0	0	0
Relapse	41 (16.3)	13 (16.4)	25 (9.2)	19 (8.0)	0	0	0	0	1 (33.3)	0
Null response	29 (11.6)	19 (10.1)	15 (5.5)	13 (5.5)	0	0	0	0	0	0
Withdrawal	62 (24.7)	-	35 (12.8)	-	1 (16.7)	-	0	-	1 (33.3)	-

Data are presented as number (%).

GT, genotype; ITT, intention-to-treat; PP, per protocol; RVR, rapid virologic response; EVR, early virologic response; pEVR, partial EVR; cEVR, complete EVR; ETR, end of treatment response; SVR, sustained virologic response.