

S3 Table. A) Total number of bacteria measured by qPCR in inocula and agar plates used on hemolytic activity of dental plaque experiments. B) p-values of Kruskal-Wallis analysis corrected for multiple comparisons. In yellow are differences that were statistically significant.

A

	Inocula	0mM	0.5mM	5mM	50mM
Patient 1	2.26x10⁷ ±3.28x10 ⁵	7.14x10¹⁰ ±1.18x10 ⁹	7.23x10¹⁰ ±8.68x10 ⁸	7.42x10¹⁰ ±3.28x10 ⁸	7.59x10¹⁰ ±1.188x10 ⁹
Patient 2	3.07x10⁷ ±4.03x10 ⁶	7.48x10¹⁰ ±8.68x10 ⁸	9.67x10¹⁰ ±6.56x10 ⁸	9.90x10¹⁰ ±3.13x10 ⁹	7.61x10¹⁰ ±4.29x10 ⁹
Patient 3	2.95x10⁷ ±1.18x10 ⁶	4.65x10¹⁰ ±8.68x10 ⁸	4.24x10¹⁰ ±8.68x10 ⁸	4.98x10¹⁰ ±4.02x10 ⁸	2.67x10¹⁰ ±1.50x10 ⁹

B

	Inocula	0mM	0.5mM	5mM	50mM
Inocula	1	0.0035	0.0000	0.0002	0.0000
0mM		1	0.0866	0.2984	0.0592
0.5mM			1	0.4234	0.1567
5mM				1	0.0674
50mM					1